

Electron driven reactions in doped He droplets

S. Denifl¹, F. Ferreira da Silva¹, S. Jaksch¹, H. Schöbel¹, F. Zappa¹, M. Dampc², O. Echt³, E. Illenberger⁴, D.K. Bohme⁵, T. D. Märk¹, P. Scheier¹

¹ *Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, Austria*

² *Department of Physics of Electronic Phenomena, Gdansk University of Technology, Poland*

³ *Department of Physics, University of New Hampshire, Durham, USA*

⁴ *Institut für physikalische und theoretische Chemie, Freie Universität Berlin, Germany*

⁵ *Department of Chemistry, York University, Toronto, Canada*

paul.scheier@uibk.ac.at

Formation of positive and negative ions upon free electron collisions with doped nano-droplets of superfluid ⁴He, are studied, utilizing a two sector field mass spectrometer. Simple diatomic molecules such as H₂ and I₂ are used as dopants as well as water, halocarbons, SF₆, CCl₄, amino acids, DNA bases and fullerenes. Molecules that collide with a droplet are picked up by the droplet and in most cases move to its center. Depending on the pressure in the pick-up cell either monomers or clusters of these dopants are formed in the droplet. Positive-ion mass spectra are recorded and compared with results obtained in the gas phase. The high mass resolution and sensitivity of the experimental setup is a prerequisite to identify novel product ions that are stabilized by the surrounding He, e.g., SF₆⁺ and CCl₄⁺.

For many dopants also negative (cluster) ion formation has been observed. The yield of these anions is measured as a function of the electron energy [1]. The molecules chosen for our first study were the nucleobases adenine, thymine and partially methylated or deuterated thymine. The DEA to these nucleobase molecules and clusters in helium droplets (see [1]) exhibits the remarkable site selectivity that we have previously established for these biomolecules in the gas phase [2]. Core excited resonances that lead to low mass fragment anions upon electron attachment in the gas phase are almost completely quenched and instead lead to the formation of dehydrogenated closed shell anions via loss of a neutral H-atom [3]. We have extended these studies to He droplets doped with several other molecules such as CHCl₃ [4], water [5], and ammonia. A surprisingly rich chemistry, driven by electrons, is observed at ultra-low (0.37K) temperatures.

Acknowledgements: This work has been supported by the FWF, Wien, Austria and the European Commission, Brussels. F.Z. gratefully acknowledges a grant from the Brazilian agency CNPq, S.D. an APART grant from the Austrian Academy of Sciences and M.D. a grant from the European Network EIPAM.

[1] S. Denifl et al., Phys. Rev. Lett. 97 (2006) 043201

[2] S. Ptasinska et al., Angew. Chem. Int. Ed. 44 (2005) 6941

[3] S. Denifl et al., ChemPhysChem (2008) in print

[4] S. Denifl et al., J. Am. Chem. Soc. 130 (2008) 5065

[5] F. Zappa et al., J. Am. Chem. Soc. 130 (2008) 5573