

Production of aluminium cluster polyanions

N. Walsh, F. Martinez, G. Marx, L. Schweikhard

Institute for physics, Ernst-Moritz-Arndt University, Greifswald, Germany

lschweik@physik.uni-greifswald.de

Interest in the physics and chemistry of multiply-charged anions has grown considerably during the last two decades. What started with a number of experimental observations of gas-phase dianionic clusters[1] and fullerenes[2,3] in the mid 1980's and early 1990's has since developed into an independent field of research investigating both the production and properties of polyanionic species[4,5,6]. One production method, the "electron-bath" technique[6] involves the exposure of stored cluster monoanions to a 'bath' of low-energy electrons for extended reaction periods. The technique, originally applied to monovalent elements[7] (Cu, Ag and Au) and later to fullerenes[8], has also proven to be suitable for the production of metal cluster polyanions composed of trivalent aluminium[9]. In addition to dianionic species[9], recently tri- (Fig.1(a)) and tetra-anionic (Fig.1(b)) aluminium clusters have been produced. This contribution presents the results of experiments in which both the production procedure and the size-dependence of aluminium cluster polyanion creation have been investigated.




Figure 1: Abundance spectra for (a) Al₁₂₅ and (b) Al₂₆₀ after exposure of the monoanions to an "electron bath".

- [1] K. Leiter et al., Int. J. Mass Spec. and Ion Process., 68(1986)341.
- [2] P.A. Limbach et al., J. Am. Chem. Soc., 113(1991)6795.
- [3] R.L. Hettich et al., Phys. Rev. Lett., 67(1991)1242.
- [4] O. Hampe et al., Chem. Phys. Lett., 354(2002)303.
- [5] B. Liu et al., Phys. Rev. Lett., 92(2004)168301.
- [6] A. Herlert et al., Phys. Scripta, T80(1999)200.
- [7] A. Herlert et al., Int. J. Mass Spec., 229(2003)19.
- [8] A. Lassesson et al., Eur. Phys. J. D, 34(2005)73.
- [9] N. Walsh et al., Eur. Phys. J. D, 43(2007)41.